arXiv:1204.3738v2 [nlin.SI] 3 Sep 2012

The Two-Component Camassa-Holm Equations CH(2,1) and CH(2,2): First-Order Integrating Factors and Conservation Laws

Marianna Euler [1], *Norbert Euler* [1] *and Thomas Wolf* [2]

[1] *Department of Engineering Sciences and Mathematics*
Luleå University of Technology
SE-971 87 Luleå, Sweden
Emails: Marianna.Euler@ltu.se; Norbert.Euler@ltu.se
[2] *Department of Mathematics, Brock University*
500 Glenridge Avenue, St.Catharines, Ontario, Canada L2S 3A1
twolf@brocku.ca

Abstract:
Recently, Holm and Ivanov, proposed and studied a class of multi-component generalisations of the Camassa-Holm equations [D D Holm and R I Ivanov, Multi-component generalizations of the CH equation: geometrical aspects, peakons and numerical examples, *J. Phys A: Math. Theor* **43**, 492001 (20pp), 2010]. We consider two of those systems, denoted by Holm and Ivanov by CH(2,1) and CH(2,2), and report a class of integrating factors and its corresponding conservation laws for these two systems. In particular, we obtain the complete sent of first-order integrating factors for the systems in Cauchy-Kovalevskaya form and evaluate the corresponding sets of conservation laws for CH(2,1) and CH(2,2).

1 Introduction

It is well known that certain conservation laws of shallow water wave equations, such as the Camassa-Holm equation [4] and the the Degasperis-Procesi equation [8], are useful to prove blow-up, cf. the papers [5], [18] and [15]. Furthermore, conservation laws play a central role in the prove of the global existence (in time) for solutions evolving from certain initial data, cf. the paper [6], and for proving the stability of peakons for both model equations, cf. the papers [7], [12] and [13]. In the context of the Camassa-Holm equation they are instrumental in the set-up of a theory of global weak solutions for nonlinear nonlocal conservation laws, cf. the considerations in the papers [2], [3] and [10]

In the current paper we derive all first-order integrating factors and its corresponding conservation laws for some recently proposed multi-component generalizations of the Camassa-Holm equation [11]. We concentrate on two explicit systems, namely CH(2,1) and CH(2,2), proposed by Holm and Ivanov in [11] (see 1.1a) – (1.1b) and (1.6a) – (1.6b) below).

We recently reported in [9] the complete set of first-order integrating factors and conservation laws for a classs of Camassa-Holm type equations, which includes the Camassa-Holm equation [4] and the the Degasperis-Procesi equation [8]. Our approach applied in this paper is based on the direct method described by Anco and Bluman in their paper [1], which can be applied to derive conservation laws of evolution equations that are in Cauchy-Kovalevskaya form. We also refer the reader to [16] and [17] for more details and alternate methods for computing conservation laws for partial differential equations and systems.

Consider the two-component Camassa-Holm equations introduced and denoted by Holm and Ivanov [11] as CH(2,1), which has the following form:

$$\sigma_1 q_t + 2qu_x + uq_x + \sigma\rho\rho_x = 0 \tag{1.1a}$$

$$\rho_t + \rho u_x + u\rho_x = 0, \tag{1.1b}$$

where

$$q = \sigma_1 u - u_{xx} + s \tag{1.2}$$

and s, σ and σ_1 are arbitrary constants. The physically interesting cases are $\sigma = \pm 1$ and $\sigma_1 = 1$ or $\sigma_1 = 0$. By defining the new dependent variables

$$u := U_1,\ u_x := U_2 \tag{1.3a}$$

$$u_{xx} := U_3,\ \rho := U_4 \tag{1.3b}$$

and the change of independent variables,

$$X := t,\ T := x, \tag{1.4}$$

we can write system (1.1a) – (1.1b) in the following Cauchy-Kovalevskaya form:

$$E_1 := U_{1,T} - U_2 = 0 \tag{1.5a}$$

$$E_2 := U_{2,T} - U_3 = 0 \tag{1.5b}$$

$$\begin{aligned} E_3 := {} & U_{3,T} - \sigma_1^2 U_1^{-1} U_{1,X} + \sigma_1 U_1^{-1} U_{3,X} - 3\sigma_1 U_2 + 2U_1^{-1} U_2 U_3 + \sigma U_1^{-2} U_4 U_{4,X} \\ & + \sigma U_1^{-2} U_2 U_4^2 - 2sU_1^{-1} U_2 = 0 \end{aligned} \tag{1.5c}$$

$$E_4 := U_{4,T} + U_1^{-1} U_{4,X} + U_1^{-1} U_2 U_4 = 0. \tag{1.5d}$$

The second 2-component Camassa-Holm equation that we study in the current paper, denoted by CH(2,2), has the form [11]

$$q_{1,t} + u_0 q_{1,x} + 2q_1 u_{0,x} + u_1 q_{2,x} + 2q_2 u_{1,x} = 0 \tag{1.6a}$$

$$q_{2,t} + u_0 q_{2,x} + 2q_2 u_{0,x} = 0, \tag{1.6b}$$

where

$$q_1 = u_1 - u_{1,xx} + s_1 \tag{1.7a}$$

$$q_2 = u_0 - u_{0,xx} + 3u_1^2 - u_{1x}^2 - 2u_1 u_{1,xx} + 4s_1 u_1 + s_2. \tag{1.7b}$$

Here $s_1,\ s_2$ are arbitrary constants. By defining the new dependent variables

$$u_0 := U_1,\ u_{0,x} := U_2,\ u_{0,xx} := U_3 \tag{1.8a}$$

$$u_1 := U_4,\ u_{1,x} := U_5,\ u_{1,xx} := U_6 \tag{1.8b}$$

and the change of independent variables (1.4), we can present (1.6a) – (1.6b) in the following Cauchy-Kovalevskaya form:

$$E_1 := U_{1,T} - U_2 = 0 \tag{1.9a}$$

$$E_2 := U_{2,T} - U_3 = 0 \tag{1.9b}$$

$$\begin{aligned}
E_3 := {} & U_{3,T} + 12U_1^{-1}U_4^3U_5 - 4U_1^{-1}U_4U_{4,X} + 2U_1^{-1}U_5U_{5,X} - 4s_1U_1^{-1}U_{4,X} \\
& +4U_5U_6 - 4s_1U_5 + 2U_1^{-1}U_2U_3 - 6U_1^{-1}U_2U_4^2 + 2U_1^{-1}U_2U_5^2 - 2s_2U_1^{-1}U_2 \\
& -4s_1U_1^{-1}U_2U_4 - 12U_1^{-2}U_2U_4^4 + 2U_1^{-1}U_6U_{4,X} - 8U_1^{-1}U_4^2U_5U_6 \\
& +16s_1U_1^{-1}U_4^2U_5 + 4U_1^{-2}U_4^2U_6U_{4,X} - 8s_1U_1^{-2}U_4^2U_{4,X} + 4U_1^{-2}U_4^2U_2U_3 \\
& +4U_1^{-2}U_4^2U_2U_5^2 + 8U_1^{-2}U_2U_4^3U_6 - 16s_1U_1^{-2}U_2U_4^3 - 4s_2U_1^{-2}U_2U_4^2 \\
& +4U_1^{-2}U_4^2U_5U_{5,X} - 4U_1^{-1}U_3U_4U_5 + 4s_2U_1^{-1}U_4U_5 - 12U_1^{-2}U_4^3U_{4,X} \\
& +2U_1^{-2}U_4^2U_{3,X} + 4U_1^{-2}U_4^3U_{6,X} - 4U_1^{-1}U_4U_5^3 - 2U_1^{-2}U_4^2U_{1,X} \\
& -U_1^{-1}U_{1,X} + U_1^{-1}U_{3,X} - 3U_2 = 0
\end{aligned} \tag{1.9c}$$

$$E_4 := U_{4,T} - U_5 = 0 \tag{1.9d}$$

$$E_5 := U_{5,T} - U_6 = 0 \tag{1.9e}$$

$$\begin{aligned}
E_6 := {} & U_{6,T} + 4U_1^{-1}U_4U_5U_6 - 8s_1U_1^{-1}U_4U_5 + 2U_1^{-1}U_5^3 - 3U_5 - U_1^{-1}U_{4,X} \\
& +U_1^{-1}U_{6,X} - 2U_1^{-2}U_4^2U_{6,X} + 6U_1^{-2}U_2U_4^3 - U_1^{-2}U_4U_{3,X} + U_1^{-2}U_4U_{1,X} \\
& +6U_1^{-2}U_4^2U_{4,X} - 2U_1^{-2}U_4U_6U_{4,X} + 4s_1U_1^{-2}U_4U_{4,X} - 2U_1^{-2}U_2U_3U_4 \\
& -2U_1^{-2}U_2U_4U_5^2 - 4U_1^{-2}U_2U_4^2U_6 + 8s_1U_1^{-2}U_2U_4^2 + 2s_2U_1^{-2}U_2U_4 \\
& -2U_1^{-2}U_4U_5U_{5,X} + 2U_1^{-1}U_3U_5 - 2s_2U_1^{-1}U_5 + 2U_1^{-1}U_2U_6 \\
& -2s_1U_1^{-1}U_2 - 6U_1^{-1}U_4^2U_5 = 0.
\end{aligned} \tag{1.9f}$$

The above first-order Cauchy-Kovalevskaya systems can now be investigated for integrating factors to derive conservation laws for the systems; which then leads to conservation laws of the systems CH(1,1) and CH(2,2) in the original variables.

2 General description

In this section we breifly describe the direct method [1] of integrating factors (or multipliers) for the general first-order Cauchy-Kovalevskaya system of six equations:

$$E_j := U_{j,T} - F_j(U_1, \dots, U_6, U_{1,X}, \dots, U_{6,X}) = 0, \quad j = 1, 2, \dots, 6. \tag{2.1}$$

Every conserved density, Φ^T, and conserved flux, Φ^X, of system (2.1) must satisfy

$$D_T \Phi^T + D_X \Phi^X \Big|_{\vec{E}=\vec{0}} = 0, \tag{2.2}$$

where, in general, both Φ^T and Φ^X are functions of X, T, U_j as well as X-derivatives of U_j. Moreover, every Φ^T requires six integrating factors, $\{\Lambda_1,\ \Lambda_2, \dots, \Lambda_6\}$, which are directly related to the conserved density by the relation [1]

$$\Lambda_k = \hat{E}[U_k]\Phi^T, \qquad k = 1, 2, \dots, 6. \tag{2.3}$$

Here $\hat{E}$ is the Euler Operator,

$$\hat{E}[U_k] := \frac{\partial}{\partial U_k} - D_T \circ \frac{\partial}{\partial U_{k,T}} + \sum_{j=1}^{q} (-1)^j D_X^j \circ \frac{\partial}{\partial U_{k,jX}}, \tag{2.4}$$

where we use the notation

$$U_{k,jX} := \frac{\partial^j U_k}{\partial X^j}.$$

The conditions on the integrating factors, $\{\Lambda_j\}$, of system (2.1) are

$$\hat{E}[U_k]\,(\Lambda_1\, E_1 + \Lambda_2\, E_2 + \cdots + \Lambda_6\, E_6) = 0, \qquad k = 1, 2, \dots, 6. \tag{2.5}$$

However, since all integrating factors of system (2.1) are adjoint symmetries of the system (2.1), we can calculate $\{\Lambda_j\}$ by the condition

$$\begin{pmatrix} L^*_{E_1}[U_1] & L^*_{E_2}[U_1] & \cdots & L^*_{E_6}[U_1] \\ L^*_{E_1}[U_2] & L^*_{E_2}[U_2] & \cdots & L^*_{E_6}[U_2] \\ \vdots & \vdots & \vdots & \vdots \\ L^*_{E_1}[U_6] & L^*_{E_2}[U_6] & \cdots & L^*_{E_6}[U_6] \end{pmatrix} \begin{pmatrix} \Lambda_1 \\ \Lambda_2 \\ \vdots \\ \Lambda_6 \end{pmatrix} \Bigg|_{\vec{E}=\vec{0}} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \tag{2.6}$$

and then require the self-adjointness condition on $\{\Lambda_j\}$ (as integrating factors are variational quatities), namely

$$\begin{pmatrix} L_{\Lambda_1}[U_1] & L_{\Lambda_1}[U_2] & \cdots & L_{\Lambda_1}[U_6] \\ L_{\Lambda_2}[U_1] & L_{\Lambda_2}[U_2] & \cdots & L_{\Lambda_2}[U_6] \\ \vdots & \vdots & \vdots & \vdots \\ L_{\Lambda_6}[U_1] & L_{\Lambda_6}[U_2] & \cdots & L_{\Lambda_6}[U_6] \end{pmatrix} \begin{pmatrix} E_1 \\ E_2 \\ \vdots \\ E_6 \end{pmatrix}$$

$$= \begin{pmatrix} L^*_{\Lambda_1}[U_1] & L^*_{\Lambda_2}[U_1] & \cdots & L^*_{\Lambda_6}[U_1] \\ L^*_{\Lambda_1}[U_2] & L^*_{\Lambda_2}[U_2] & \cdots & L^*_{\Lambda_6}[U_2] \\ \vdots & \vdots & \vdots & \vdots \\ L^*_{\Lambda_1}[U_6] & L^*_{\Lambda_2}[U_6] & \cdots & L^*_{\Lambda_6}[U_6] \end{pmatrix} \begin{pmatrix} E_1 \\ E_2 \\ \vdots \\ E_6 \end{pmatrix}. \qquad (2.7)$$

$$(2.8)$$

Here L is the linear operator and L^* its adjoint:

$$L_P[U_j] := \frac{\partial P}{\partial U_j} + \sum_{i=1}^{p} \frac{\partial P}{\partial U_{j,iT}}\ D_T^i + \sum_{k=1}^{q} \frac{\partial P}{\partial U_{j,kX}}\ D_X^k \qquad (2.9a)$$

$$L_P^*[U_j] := \frac{\partial P}{\partial U_j} + \sum_{i=1}^{p} (-1)^i\ D_T^i \circ \frac{\partial P}{\partial U_{j,iT}} + \sum_{k=1}^{q} (-1)^k\ D_X^k \circ \frac{\partial P}{\partial U_{j,kX}}. \qquad (2.9b)$$

Note that the self-adjointness condition, (2.7), is independent of the form of the evolution system (2.1) and only depends on the functional arguments of $\{\Lambda_j\}$ as well as the number of equations in the system.

3 Integrating factors for system (1.5a) – (1.5d) and conservation laws for (1.1a) – (1.1b):

Solving conditions (2.6) and (2.7) for system (1.5a) – (1.5d), the complete set of first-order integrating factors $\{\Lambda_1, \ldots, \Lambda_4\}$, of the form

$$\Lambda_j = \Lambda_j(X, T, U_1, \ldots, U_4, U_{1,X}, \ldots, U_{4,X}), \quad j = 1, 2, \ldots, 4$$

for arbitrary σ, $\sigma_1 \neq 0$ and s is as follows:

$$\Lambda_1 = \lambda_1 U_4 + 2\lambda_2 \left(s + \frac{3}{2}\sigma_1 U_1 - \frac{1}{2}U_3 \right)$$
$$-\lambda_3 \left(\sigma_1 U_{2,X} - 2sU_1 - \sigma U_4^2 - 3\sigma_1 U_1^2 + 2U_1 U_3 \right) \sigma_1^{-1} \tag{3.1a}$$

$$\Lambda_2 = -\lambda_2 U_2 + \lambda_3 U_{1,X} \tag{3.1b}$$

$$\Lambda_3 = -\lambda_2 U_1 - \lambda_3 \sigma_1^{-1} U_1^2 \tag{3.1c}$$

$$\Lambda_4 = \lambda_1 U_1 + \lambda_2 \sigma U_4 + 2\lambda_3 \sigma \sigma_1^{-1} U_1 U_4 \tag{3.1d}$$

where λ_j are arbitrary constants. This leads to the following three sets of conserved density, Φ^t, and conserved flux, Φ^x, for the original system (1.1a) – (1.1b) (separated by means of the arbitrary λ_1, λ_2 and λ_3, respectively):

$$\Phi_1^t = \rho \tag{3.2a}$$

$$\Phi_1^x = u\rho \tag{3.2b}$$

$$\Phi_2^t = \sigma_1^2 u - \sigma_1 u_{xx} \tag{3.3a}$$

$$\Phi_2^x = 2su + \frac{1}{2}\sigma\rho^2 + \frac{3}{2}\sigma_1 u^2 - uu_{xx} - \frac{1}{2}u_x^2 \tag{3.3b}$$

$$\Phi_3^t = \left(-\sigma_1 u_x^2 + \frac{1}{2}\sigma\rho^2 + \frac{1}{2}\sigma_1^2 u^2 - \sigma_1 uu_{xx} + \frac{1}{2}\sigma_1 u_x^2 \right) \sigma_1^{-1} \tag{3.4a}$$

$$\Phi_3^x = \left(\sigma_1 u_x u_t + su_x^2 + \sigma u\rho^2 + \sigma_1 u^3 - u^2 u_{xx} \right) \sigma_1^{-1} \tag{3.4b}$$

Some special must be considered:

Special Case 1: $\sigma = 0$ with σ_1 arbitrary, but nonzero, and s arbitrary. The integrating factors are as follows:

$$\Lambda_1 = \frac{2U_3 - 3\sigma_1 U_1 - 2s}{(\sigma_1 U_1 - U_3 + s)^{1/2}}, \quad \Lambda_2 = 0 \tag{3.5}$$

$$\Lambda_3 = \frac{U_1}{(\sigma_1 U_1 - U_3 + s)^{1/2}}, \quad \Lambda_4 = 0. \tag{3.6}$$

and the corresponding conserved current for system (1.5a) – (1.5d) is

$$\Phi^t = \sigma_1 \left(\sigma_1 u - u_{xx} + s \right)^{1/2} \tag{3.7a}$$

$$\Phi^x = \left(\sigma_1 u - u_{xx} + s \right)^{1/2}. \tag{3.7b}$$

Special Case 2: $\sigma = 0$ with $\sigma_1 = 1$ and s arbitrary. The integrating factors are as follows:

$$\Lambda_1 = -\frac{U_1 W^3 H'(W)}{U_4} + 2U_4 H(W), \quad \Lambda_2 = 0 \tag{3.8a}$$

$$\Lambda_3 = \frac{U_1 W^3 H'(W)}{U_4}, \quad \Lambda_4 = 2U_1 \left(W H'(W) + H(W) \right), \tag{3.8b}$$

where $H(W)$ is an arbitrary differentiable function with

$$W := \frac{U_4}{(U_1 - U_3 + s)^{1/2}}. \tag{3.9}$$

The conserved current for system (1.5a) – (1.5d) is then

$$\Phi^t = H(w)\rho \tag{3.10a}$$

$$\Phi^x = H(w)u\rho. \tag{3.10b}$$

Here the argument, w, in the arbitrary function H, is

$$w := \frac{\rho}{(u - u_{xx} + s)^{1/2}}. \tag{3.11}$$

Special Case 3: σ arbitrary, but nonzero, with $\sigma_1 = 1$ and s arbitrary. The integrating factors are as follows:

$$\Lambda_1 = \frac{U_3 - 2U_1 - s}{\sigma U_4}, \quad \Lambda_2 = 0, \tag{3.12a}$$

$$\Lambda_3 = \frac{U_1}{\sigma U_4}, \quad \Lambda_4 = \frac{s U_1 - \sigma U_4^2 + U_1^2 - U_1 U_3}{\sigma U_4^2}. \tag{3.12b}$$

The conserved current for system (1.5a) – (1.5d) is then

$$\Phi^t = \frac{u_{xx} - u - s}{\sigma\rho} \tag{3.13a}$$

$$\Phi^x = \frac{u u_{xx} - u^2 - \sigma\rho^2 - s u}{\sigma\rho}. \tag{3.13b}$$

Special Case 4: $\sigma_1 = 0$ with σ and s arbitrary. The integrating factors are as follows:

$$\Lambda_1 = (U_3 - 2s) H(X, W), \quad \Lambda_2 = U_2 H(X, W) \tag{3.14a}$$

$$\Lambda_3 = U_1 H(X, W), \quad \Lambda_4 = -\sigma U_4 H(X, W), \tag{3.14b}$$

where H is an arbitrary differentiable function and

$$W := \frac{1}{2} \left(-4 s U_1 - \sigma U_4^2 + 2 U_1 U_3 + U_2^2 \right). \tag{3.15}$$

The conserved current for system (1.5a) – (1.5d) is then

$$\Phi^t = 0 \tag{3.16a}$$

$$\Phi^x = wH(t,w), \tag{3.16b}$$

where

$$w := \frac{1}{2}\left(-4su - \sigma\rho^2 + 2uu_{xx} + u_x^2\right). \tag{3.17}$$

Special Case 5: $\sigma_1 = 0$ with σ arbitrary, but nonzero, and s arbitrary. The integrating factors are as follows:

$$\Lambda_1 = \frac{2}{\sigma}\left(sU_1 + \frac{\sigma}{2}U_4^2 - U_1U_3\right), \quad \Lambda_2 = 0 \tag{3.18a}$$

$$\Lambda_3 = -\frac{U_1^2}{\sigma}, \quad \Lambda_4 = 2U_1U_4. \tag{3.18b}$$

The conserved current for system (1.5a) – (1.5d) is then

$$\Phi^t = \frac{\rho^2}{2} \tag{3.19a}$$

$$\Phi^x = \frac{1}{\sigma}\left(su^2 + \sigma u\rho^2 - u^2u_{xx}\right). \tag{3.19b}$$

4 Integrating factors for system (1.9a) – (1.9f) and conservation laws for (1.6a) – (1.6b):

Solving conditions (2.6) and (2.7) for system (1.9a) – (1.9f), the complete set of first-order integrating factors $\{\Lambda_1, \ldots, \Lambda_6\}$, of the form

$$\Lambda_j = \Lambda_j(X, T, U_1, \ldots, U_6, U_{1,X}, \ldots, U_{6,X}), \quad j = 1, 2, \ldots, 6$$

are the following:

$$\begin{aligned}
\Lambda_1 ={}& \lambda_1 \left(2U_6U_1 + 2U_3U_4 + 2U_4U_5^2 + 4U_4^2U_6 - 6U_4^3 - 2s_1U_1 - 8s_1U_4^2 - 6U_1U_4\right.\\
&\quad \left.-2s_2U_4 + U_{5,X}\right) + \lambda_2 \left(U_3 + 2U_5^2 - 4s_1U_4 - 3U_1 - 2s_2\right)\\
&\quad +\lambda_3 \left(U_6 - 3U_4 - 2s_1\right) + \lambda_4 \left(-8s_1^2U_4 - 2s_1s_2 - s_1U_1 + 2s_1U_3 - 22s_1U_4^2\right.\\
&+12s_1U_4U_6 + 2s_1U_5^2 - 4s_2U_4 + 2s_2U_6 - 3U_1U_4 + U_1U_6 + 4U_3U_4 - 2U_3U_6 - 12U_4^3\\
&\left.+14U_4^2U_6 + 4U_4U_5^2 - 4U_4U_6^2 - 2U_5U_6\right) Z^{-3/2}\\
&+\frac{\lambda_5}{2} \left(-8s_1U_4 - 2s_2 - 3U_1 + 2U_3 - 6U_4^2 + 4U_4U_6 + 2U_5^2\right) Z^{-1/2}
\end{aligned}\tag{4.1a}$$

$$\Lambda_2 = -\lambda_1U_{4,X} + \lambda_2U_2 + \lambda_3U_5 \tag{4.1b}$$

$$\begin{aligned}
\Lambda_3 ={}& 2\lambda_1U_1U_4 + \lambda_2 \left(U_1 - 2U_4^2\right) + \lambda_3U_4\\
&+\lambda_4 \left(-s_1U_1 + 8s_1U_4^2 + 2s_2U_4 + U_1U_4 + U_1U_6 - 2U_3U_4 + 6U_4^3\right.\\
&\left.-4U_4^2U_6 - 2U_4U_5^2\right) Z^{-3/2} + \frac{\lambda_5}{2}U_1Z^{-1/2}
\end{aligned}\tag{4.1c}$$

$$\begin{aligned}
\Lambda_4 ={}& \lambda_1 \left(2U_1U_5^2 + 2U_1U_3 + 2U_4U_{5,X} - 2s_2U_1 - 3U_1^2 - 18U_1U_4^2 + U_{2,X}\right.\\
&\quad \left.-16s_1U_1U_4 + 8U_1U_4U_6\right) + \lambda_2 \left(24U_4^3 - 4U_3U_4 - 4U_4U_5^2 - 12U_4^2U_6 - 2U_{5,X}\right.\\
&\quad \left.+24s_1U_4^2 - 4s_1U_1 + 4s_2U_4\right) + \lambda_3 \left(U_3 + 4U_4U_6 - 3U_1 + 2U_5^2 - 12U_4^2\right.\\
&\quad \left.-12s_1U_4 - 2s_2\right) + 2\lambda_4 \left(2s_1^2U_1 - 48s_1^2U_4^2 - 20s_1s_2U_4 - 19s_1U_1U_4 - 3s_1U_1U_6\right.\\
&+20s_1U_3U_4 - 84s_1U_4^3 + 48s_1U_4^2U_6 + 20s_1U_4U_5^2 - 2s_2^2 - 5s_2U_1 + 4s_2U_3 - 18s_2U_4^2\\
&+10s_2U_4U_6 + 4s_2U_5^2 - 3U_1^2 + 5U_1U_3 - 18U_1U_4^2 + 8U_1U_4U_6 + 5U_1U_5^2 + U_1U_6^2 - 2U_3^2\\
&+18U_3U_4^2 - 10U_3U_4U_6 - 4U_3U_5^2 - 36U_4^4 + 42U_4^3U_6 + 18U_4^2U_5^2 - 12U_4^2U_6^2\\
&\left.-10U_4U_5^2U_6 - 2U_5^4\right) Z^{-3/2} + \lambda_5U_1 \left(-2s_1 + 3U_4 + U_6\right) Z^{-1/2}
\end{aligned}\tag{4.1d}$$

$$\begin{aligned}
\Lambda_5 ={}& \lambda_1 \left(4U_1U_4U_5 - U_{1,X} - 2U_4U_{4,X}\right) + \lambda_2 \left(4U_1U_5 - 4U_4^2U_5 + 2U_{4,X}\right)\\
&\quad +\lambda_3 \left(U_2 + 4U_4U_5\right) - 2\lambda_4U_5 \left(s_1U_1 - 8s_1U_4^2 - 2s_2U_4 - U_1U_4 - U_1U_6 + 2U_3U_4\right.\\
&\left.-6U_4^3 + 4U_4^2U_6 + 2U_4U_5^2\right) Z^{-3/2} + \lambda_5U_1U_5Z^{-1/2}
\end{aligned}\tag{4.1e}$$

$$\begin{aligned}
\Lambda_6 ={}& \lambda_1 \left(U_1^2 + 4U_1U_4^2\right) - 4\lambda_2U_4^3 + \lambda_3 \left(U_1 + 2U_4^2\right) + 2\lambda_4 \left(3s_1U_1U_4 + 8s_1U_4^3 + s_2U_1\right.\\
&+2s_2U_4^2 - U_1^2 - U_1U_3 + 4U_1U_4^2 - U_1U_4U_6 - U_1U_5^2 - 2U_3U_4^2 + 6U_4^4\\
&\left.-4U_4^3U_6 - 2U_4^2U_5^2\right) Z^{-3/2} + \lambda_5U_1U_4Z^{-1/2},
\end{aligned}\tag{4.1f}$$

where

$$Z := s_1U_4 - s_2 - U_1 + U_3 - 3U_4^2 + 2U_4U_6 + U_5^2. \tag{4.2}$$

This leads to the following set of three conserved densities and conserved flux for the system (1.6a) – (1.6b):

$$\Phi_1^t = u_1u_{0,xx} + u_1^2u_{1,xx} - u_0u_1 - 2s_1u_1^2 - 2u_1^3 \tag{4.3a}$$

$$\begin{aligned}\Phi_1^x = &\left(u_0 + u_1^2\right) u_{1,xt} + 2u_0u_1u_{0,xx} + 2u_0u_1u_{1,x}^2 + \left(4u_0u_1^2 + u_0^2\right) u_{1,xx} \\ &-\frac{1}{2}u_0^2\left(6u_1 + 2s_1\right) - u_0\left(6u_1^3 + 2s_2u_1 + 8s_1u_1^2\right) - u_{0,x}u_{1,t}\end{aligned} \tag{4.3b}$$

$$\Phi_2^t = 2u_1u_{1,xx} + u_{0,xx} - u_0 - 2u_1^2 + 2u_{1,x}^2 - 4s_1u_1 \tag{4.4a}$$

$$\begin{aligned}\Phi_2^x = &-2u_1u_{1,xt} + (u_0 - 2u_1^2)u_{0,xx} - 4u_1^3u_{1,xx} + \frac{1}{2}u_{0,x}^2 + 2(u_0 - u_1^2)u_{1,x}^2 \\ &-2(s_2 + 2s_1u_1)u_0 - \frac{3}{2}u_0^2 + 2u_1^2(s_2 + 4s_1u_1 + 3u_1^2)\end{aligned} \tag{4.4b}$$

$$\Phi_3^t = u_{1,xx} - u_1 \tag{4.5a}$$

$$\begin{aligned}\Phi_3^x = &(u_0 + 2u_1^2)u_{1,xx} + u_1u_{0,xx} + u_{0,x}u_{1,x} + 2u_1u_{1,x}^2 - (2s_1 + 3u_1)u_0 \\ &-2u_1(s_2 + 3s_1u_1 + 2u_1^2).\end{aligned} \tag{4.5b}$$

$$\Phi_4^t = 2\left(s_1 - u_{1,xx} + u_1\right) z^{-1/2} \tag{4.6a}$$

$$\begin{aligned}\Phi_4^x = &2\left(s_1u_0 + 8s_1u_1^2 + 2s_2u_1 + 3u_0u_1 - u_0u_{1,xx} - 2u_{0,xx}u_1 + 6u_1^3\right. \\ &\left.-4u_1^2u_{1,xx} - 2u_1u_{1,x}^2\right) z^{-1/2}\end{aligned} \tag{4.6b}$$

$$\Phi_5^t = z^{1/2} \tag{4.7a}$$

$$\Phi_5^x = u_0z^{1/2}, \tag{4.7b}$$

where

$$z := s_1u_1 - s_2 - u_0 + u_{0,xx} - 3u_1^2 + 2u_1u_{1,xx} + u_{1,x}^2. \tag{4.8}$$

5 Concluding remarks

We have derived the complete set of first-order integrating factors for the systems CH(2,1) and CH(2,2) in Cauchy-Kovalevskaya form. The corresponding sets of conservation laws related to these integrating factors have been derived for both these systems. It would certainly be interesting to calculate higher-order integrating factors, although the computations involved for such calculations appear to be rather challenging. We aim to report some results in a future paper.

We expect that the same method than was applied here could also be used to find conservation laws for more general CH-systems proposed in [11] and [14].

Acknowledgements

ME and NE thanks the New Jersey Institute of Technology for there hospitality during their sabbatical leave at this institute. ME and NE also thank the Wenner-Gran Foundation and Luleå University of Technology for financial support.

References

[1] C. S. Anco and G. Bluman, Direct construction method for conservation laws of partial differential equations Part II: General treatment, *Euro. Jnl of Applied Mathematics* **13**, 567–585, 2002.

[2] A. Bressan and A. Constantin, Global conservative solutions of the Camassa-Holm equation, *Arch. Ration. Mech. Anal.* **183** 215-239, 2007.

[3] A. Bressan and A. Constantin, Global dissipative solutions of the Camassa-Holm equation, *Anal. Appl.* **5**, 127, 2007.

[4] R. Camassa and D. D. Holm, An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.* **71**, 11, 1661–1664, 1993.

[5] A. Constantin, On the Cauchy problem for the periodic Camassa-Holm equation, *J. Differential Equations* **141**, 218 -235, 1997.

[6] A. Constantin and J. Escher, Well-posedness, global existence, and blowup phenomena for a periodic quasi-linear hyperbolic equation, *Comm. Pure Appl. Math.* **51**, 475 -504, 1998.

[7] A. Constantin and W. Strauss, Stability of peakons, *Comm. Pure Appl. Math.* **53**, 603- 610, 2000.

[8] A. Degasperis, D. D. Holm and A. N. W. Hone, A New Integrable Equation with Peakon Solutions, *Theor. and Math. Phys.* **133**, 2, 1463–1474, 2002.

[9] M. Euler and N. Euler, Integrating factors and conservation laws for some Camassa-Holm type equations, *Commun. Pure Appl. Anal.* **11**, 1421-1430, 2012.

[10] H. Holden and X. Raynaud, Golbal semigroup of conservative solutions of the nonlinear variational wave equation, *Arch. Ration. Mech. Anal.* **201**, 871-964, 2011.

[11] D. D. Holm and R. I. Ivanov, Multi-component generalizations of the CH equation: geometrical aspects, peakons and numerical examples, *J. Phys A: Math. Theor* **43**, 492001 (20pp), 2010.

[12] J. Lenells, A variational approach to the stability of periodic peakons, *J. Nonlinear Math. Phys.* **11**, 151 -163, 2004.;

[13] Z. Lin and Y. Liu, Stability of peakons for the Degasperis-Procesi equation, *Comm. Pure Appl. Math.* **62**, 125 -146, 2009.

[14] V. Novikov, Generalisations of the Camassa-Holm equation, *J. Phys. A* **42** 34, 342002, 14 pp, 2009.

[15] E. Wahlen, On the blow-up of solutions to the periodic Camassa-Holm equation, *NoDEA* **13**, 643-653, 2007.

[16] T. Wolf, A comparison of four approaches to the calculation of conservation laws, *Euro. Jnl of Applied Mathematics* **13** (part 2), 129 -152, 2002.

[17] T. Wolf, Applications of CRACK in the Classification of Integrable Systems, *CRM Proceedings and Lecture Notes*, **37**, 283 -300, 2004.

[18] X. Wu, On the Cauchy problem for the periodic generalized Degasperis-Procesi equation, *J. Funct. Anal.* **260**, 1428 -1445, 2011